

INSTITUTE FOR ANIMAL HEALTH

Director: Professor Martin W. Shirley, PhD

PIRBRIGHT LABORATORY

Ash Road,

Pirbright,

Surrey,

GU24 ONF

Intn Tel: 00 44 1483 232441

Tel: 01483 232441 Fax: 01483 232621

FMD Vaccine Matching Strain Differentiation Report

Lab Reference WRL Batch Number:

Sender Details:

WRLFMD/2010/00028

Date Received: Country of Origin: Date Reported: 25th June 2010

Myanmar

29th November 2010

Report no:	2dmVNT					LPBE				
Vaccine:	า	0	O Ind	0	0		0	O BFS	O Tai	0
Field Isolate:	VNT	Bfs	R2/75	Manisa	Taw98	LPBE	3039	1860	189/87	Manisa
O Mya										
13/2009	Mean	0.10	0.64	0.16	0.20	Mean	0.27	DNT	DNT	0.21
O Mya 3/2010	Mean	0.16	0.57	0.39	0.15	Mean	DNT	DNT	DNT	DNT

Results Approved By:





To help us improve the quality of our service, please send any suggestions or requests to the Reference Laboratory by fax (+44 (0) 1483 232621 or email: elizabeth.wilson@bbsrc.ac.uk)

Interpretation of Results

In the case of Virus Neutralisation Test (VNT):

 $r_1 = \ge 0.3$. Suggests that there is a close relationship between field isolate and vaccine strain. A potent vaccine containing the vaccine strain is likely to confer protection.

 r_1 = < 0.3. Suggests that the field isolate is so different from the vaccine strain that the vaccine is unlikely to protect.

ND = Not done.

In the case of Liquid Phase Blocking Elisa (LPBE):

 r_1 = 0.4-1.0. Suggests that there is a close relationship between field isolate and vaccine strain. A potent vaccine containing the vaccine strain is likely to confer protection.

 r_1 = 0.2-0.39, Suggests that the field isolate is antigenically related to the vaccine strain. The vaccine strain might be suitable for use if no closer match can be found provided that a potent vaccine is used and animals are preferably immunised more than once.

 r_1 = <0.2. Suggests that the field isolate is so different from the vaccine strain that the vaccine is unlikely to protect.

DNT = Did not trap.

ND = Not done.